




Amsterdam UMC  
Universitair Medische Centra

# Transfusion-associated circulatory overload

Dr. R. Klanderman, MD PhD  
AIOS Anesthesiologie / Intensive Care

NVB 10-05-2023







# Indicaties

- Componenten therapie:

- RBC's

↓ Hb

- Plaatjes

↓ aantal / slechte functie

- Plasma

↓ stollingsfactoren

- Alle andere effecten ongewenst!





# TACO

- Komt vaak voor
  - #1 transfusie mortaliteit NL, FR, USA
  - 1% alle transfusies<sup>1</sup> Transfusies:
    - NL: 350.000/jr
    - USA: 11.000.000/jr
  - 6% op de IC<sup>2</sup> / OK<sup>3</sup>
- Mortaliteit: 10%<sup>4</sup>

1. Roubinian *Vox Sang* 2017;  
3. Clifford *Anesthesiology* 2015;

2. Li *Transfusion* 2011;  
4. Roubinian *Crit Care Med* 2018





# Waarom TACO?

- #1 oorzaak overlijden door transfusie

## Bijzonder:

- 30 - 50% van TACO na 1 unit<sup>5</sup>
- Incidentie ↓ na leukoreductie<sup>6</sup>
- 33% koorts<sup>7</sup>




Fig. 2. The annual incidence of TACO before (1993-1999) and after (2001-2007) the introduction of universal leukoreduction (ULR) in the year 2000 is shown. The incidence rate was five or fewer cases per 100,000 components transfused in 6 of the 7 years after the introduction of ULR, compared with only 2 of the 7 years before the introduction of ULR.

5. Piccin *Transfusion* 2015

6. Blumberg *Transfusion* 2010



7. Parmar *Vox Sang* 2017




# Thesis doelen:

- 1. Is TACO meer dan volume overbelasting? Hoe ontstaat het?**
  
  
  
  
  
  
- 2. Kunnen we TACO voorkomen?**

# Diermodel




Gezonde dieren =  $\emptyset$  effect






### Myocardial Infarction



### Acute Kidney Injury



# Diermodel

- 1<sup>e</sup> TACO dier model

- Two-hit model

- RBCs slechter dan vocht

TACO is meer dan  
volume overbelasting!




# Thesis doelen:

- 1. Is TACO meer dan volume overbelasting? Hoe ontstaat het?**
  - JA!
  - Two-hit model
  
- 2. Kunnen we TACO voorkomen?**




# Snelheid vs. volume

- Transfusion volume
- Transfusion speed






# Furosemide prophylaxe





# Furosemide prophylaxe

- Direct voor transfusie
- Lage dosis  $\pm$  40mg
- Hoge dosis  $\pm$  120mg





# Thesis doelen:

1. Is TACO meer dan volume overbelasting? Hoe ontstaat het?

- JA!
- Two-hit model

2. Kunnen we TACO voorkomen?

- Transfusie: snelheid




volume

- Furosemide, 1e evidence



# Pathofysiologie






# Toekomst

- Strategie per transfusie product

- TACO - TRALI spectrum

- Individueel transfusie beleid
  - Laag volume producten
  - Snelheid
  - Profylaxe





# Met dank aan

- **Amsterdam UMC**
  - Alexander Vlaar
  - Markus Hollmann
  - Margreeth Vroom
  - Bart Geerts
  - Esther Bulle
  - Phillipa Phelps
- **Sanquin Blood Bank**
  - Dirk de Korte
  - Robin van Bruggen
  - Jaap van Buul
  - Rick Kapur



# References:



1. Roubinian NH, et al. Incidence and clinical characteristics of transfusion-associated circulatory overload using an active surveillance algorithm. *Vox Sang*. 2017 Jan;112(1)
2. Li G, et al. Incidence and transfusion risk factors for transfusion-associated circulatory overload among medical intensive care unit patients. *Transfusion* 2011;51:338-43.
3. Clifford L, et al. Characterizing the epidemiology of perioperative transfusion-associated circulatory overload. *Anesthesiology* 2015;122:21-8.
4. Roubinian NH, et al. Contemporary risk factors and outcomes of transfusion-associated circulatory overload. *Crit Care Med* 2018;46(4):577-85
5. Piccin A, et al. Transfusion-associated circulatory overload in Ireland: a review of cases reported to the National Haemovigilance Office 2000 to 2010. *Transfusion*. 2015 Jun;55(6):1223-30
6. Blumberg N, et al. An association between decreased cardiopulmonary complications (transfusion-related acute lung injury and transfusion-associated circulatory overload) and implementation of universal leukoreduction of blood transfusions. *Transfusion*. 2010 Dec;50(12):2738-44
7. Parmar N, et al. The association of fever with transfusion-associated circulatory overload. *Vox Sang*. 2017 Jan;112(1):70-78
8. Klanderman RB, et al. Volume incompliance and transfusion are essential for transfusion-associated circulatory overload: a novel animal model. *Transfusion*. 2019 Dec;59(12):3617-3627
9. Klanderman RB, et al. Differential effects of speed and volume on transfusion-associated circulatory overload: A randomized study in rats. *Vox Sang*. 2022
10. Klanderman RB, et al. Prophylactic furosemide to prevent transfusion-associated circulatory overload: a randomized controlled study in rats. *Sci Rep*. 2022 Jul 15;12(1):12127
11. Bulle EB, et al. The recipe for TACO: A narrative review on the pathophysiology and potential mitigation strategies of transfusion-associated circulatory overload. *Blood Rev*. 2022 Mar;52:100891